

Unintentional Death by Fall

Falls and fall-related injuries account for more than 20 percent of all unintentional (accidental) deaths to Idaho residents. Between 2004 and 2006, an average of 10 Idaho residents died per month as the result of a fall or fall-related injury. In 2005, the National Safety Council reported 8 million people were treated in emergency rooms across the country for fall-related injuries; making falls the leading cause of non-fatal injuries for all age groups except those aged 15-24.¹ Falls account for 1.3 percent of all deaths to Idaho residents. Between 2004 and 2006, a total of 356 Idahoans died as the result of an unintentional fall – 301 of those individuals were over the age of 64.

2004-2006 Idaho Quick Facts:

Mean age of death by fall in Idaho:

79 (All)

Median age of death by fall in Idaho:

83 (Women)

74 (Men)

79 (All)

Annual Number of Deaths Caused by a Fall:

141 (2006)

101 (2005)

114 (2004)

113 (Average)

85 percent of individuals who die from falls are aged 65 and older (2004-2006).

48 percent of falls occurred to individuals aged 85 and older.

Head (35%) and hip (33%) injuries are the most common injuries incurred in a fall.

Fall injuries are 6 times more likely to occur in a residential location than any other.

Who is Affected?

In Idaho, falls represent the greatest risk of unintentional death for people aged 65 and older. Between 2004 and 2006, an average of 113 individuals died per year as the result of a fall or fall-related injury; 100 of those deaths occurred to individuals aged 65 and older.


Between 2004 and 2006, 85 percent of all unintentional deaths by fall occurred to individuals aged 65 and older. The average age of death by fall for Idaho residents was 79.

An estimated one-third of individuals aged 65 and older fall once during a year.² The likelihood of a severe or life-threatening injury increases significantly with age.³ In addition, it is estimated that two-thirds of older adults who fall will fall again within the next 6 months.⁴


In Idaho, the three-year aggregate age and sex-specific death rates indicate that within specific age groups men are more likely to die from a fall than women. However, when taken as a whole, women aged 65 and older have a 7 percent increased risk over men for dying from a fall (63.1 and 58.9 per 100,000, respectively). In comparison, national rates are not consistent with Idaho rates.

In 2005 (latest national rates available from the Centers for Disease Control and Prevention), the national rates for unintentional injury deaths due to falls were 50.4 and 34.7 per 100,000 population, respectively, for men and women.

Idaho Resident Unintentional-Injury Death by Fall 2004-2006 by Age Group n = 356




Idaho Resident Unintentional-Injury Death by Fall 2004-2006 by Gender and Age Group



Where are Idahoans Dying from a Fall?


Idaho Resident Unintentional-Injury Death by Fall 2004-2006 by Injury Location

n = 356



Idaho Resident Unintentional-Injury Death by Fall 2004-2006 by Type of Fall in Residence

n = 307



A total of 48 (16 percent) in-home accidental fall deaths between 2004 and 2006 were caused by slipping, tripping, stumbling or some other noted loss of balance. Fourteen percent (43 individuals) involved falling out of or off of a bed, wheelchair, or some other kind of furniture. Eight percent of in-home fall deaths involved stairs.

Across Idaho, Districts 4 and 7 were the only districts whose rate varied significantly from Idaho's. District 4 had the lowest rate of fall deaths (5.9 per 100,000 population) - significantly lower than the total for the state of Idaho. District 7 reported a rate of 11.7 deaths per 100,000 - a rate significantly higher than the state death rate (8.3 per 100,000).


Idahoans are more likely to die from a fall within their own home than any other location. Eighty-six percent (307 deaths) of all fall-related deaths between 2004 and 2006 occurred within the home, an assisted living center, or another residential type of institution. Outdoor accidents were the most prevalent location for non-residential fall-related death, accounting for a 3.9 percent (14) of deaths during this time period.

Idaho Resident Unintentional-Injury Death by Fall 2004-2006 by Location

Location of Accident	Number	Percent
All Locations	356	100.0%
Residential	307	86.2%
Home	218	61.2%
Assisted Living Center, Veterans Home, or other Residential Institution	89	25.0%
Non-Residential	40	11.2%
Outdoors location: Forest, Trail, Campsite, Cliff, or Hot Springs.	14	3.9%
Supermarket, Airport, Bar, or other Trade/Service Area	11	3.1%
Hospital, School, or other Public Building	6	1.7%
Industrial, Construction, or other Worksite	5	1.4%
Street, Highway, Sidewalk, or Parking Lot	4	1.1%
Sports Complex, Athletic Venue or Arena	-	-
Farm	-	-
Unknown /Unspecified	9	2.5%

Unintentional fall deaths in assisted living centers, nursing homes, and other residential institutions accounted for 25 percent of all fall deaths to Idaho residents. It is estimated that 60 percent of nursing home residents fall each year in the United States.⁶


Idaho Resident Unintentional-Injury Death by Fall 2004-2006 by District of Residence



How are Idahoans Accidentally Dying?

In Idaho, falls are the second leading cause of accidental death for all age groups, and the leading cause of accidental death for individuals aged 65 and older. Falls and fall-related injuries account for more deaths in this age group than motor vehicle accidents, choking, poisoning, exposure to fire, and drowning combined.

Idaho Resident Unintentional-Injury Fall Death 2004-2006 by Primary Injury n = 356




The majority of fall-related deaths were caused by head trauma (35 percent) and hip fractures (33 percent).

Of injuries sustained in a fall, head trauma and hip fractures are particularly dangerous, proving nationally to be a serious threat to individual independence and quality of life.⁵

Half of unintentional fall deaths and 8 percent of hospitalizations for non-fatal falls are the result of a traumatic brain injury.⁸ These injuries often result in long-term cognitive, emotional, and functional impairments.⁸

Across the country, more than 90 percent of hip fractures among adults aged 65 and older are caused by falls.⁷ The CDC estimates of those older adults living independently before their hip fracture, only half were able to live independently a year later.³

Idaho Resident Unintentional-Injury Deaths 2004-2006 by Cause and Age Group




Note: Accidental poisoning deaths include accidental poisonings by drugs and other biological substances; unspecified solid or liquid substances; and exposure to poisonous gasses and vapors.

Trend

The National Safety Council estimates that in 2005 there were 41.9 million visits to hospitals across the country as a result of accidental injuries.¹ Of those, approximately 21 percent were the result of a fall. From 1985 to 2006, the rate of fall-related death increased 76.9 percent in the U.S. In Idaho, the rate increased 92.0 percent over the same time period.


Unintentional-Injury Death by Fall 1985-2006 Idaho and U.S.



U.S. 2006 rate is preliminary. Statistics for 1985-1998 are based on ICD-9 Codes 880-888 and were modified with the comparability ratio of 0.772; statistics for 1999-2006 are based on ICD-10 Codes W00.0-W19.9.

Risk Factors and Conditions Associated with Falls

Percent of Idaho Adults Who have Experienced a Fall or Fall-Related Injury by Condition, 2005 BRFSS




Overweight defined as those individuals with a Body Mass Index (BMI) greater than or equal to 25. Obese defined as those individuals with a BMI greater than or equal to 30. * Percent is significantly higher for adults with condition than adults without the condition at the 95% confidence interval.

2005 data from the Behavior Risk Factor Surveillance System (BRFSS) showed that Idahoans diagnosed with a stroke, heart attack, heart disease, diabetes, or arthritis were significantly more likely to have experienced an injury from a fall in the past 12 months than those who did not have these conditions.

The strongest correlation was found between the use of special equipment (wheelchair, cane, special bed, or special telephone) and the likelihood of having experienced a fall. Individuals who regularly use special equipment are more than twice as likely to experience a fall (43.9 percent versus 17.3 percent) and more than three times as likely to experience an injury due to a fall than those who do not use special equipment (24.7 percent and 7.6 percent, respectively). It is important to note that correlation between falls and the use of special equipment does not necessarily imply causation.

Percent of Idaho Adults Who Have Experienced a Fall or Fall-Related Injury by Use of Special Equipment, 2005 BRFSS



Idaho Department of Health and Welfare, Bureau of Vital Records and Health Statistics (August 2008)
 Costs associated with this fact sheet are available from Idaho Department of Health and Welfare. HW-1206

- National Safety Council, Injury Facts, 2008 Edition.
- Hausdorff JM, Rios DA, Edelber HK. Gait variability and fall risk in community-living older adults: a 1-year prospective study. Archives of Physical Medicine and Rehabilitation 2001;82(8):1050-6.
- Sleet, DA., Moffett, DB., Stevens, JA. (2008) CDC's research portfolio in older adult fall prevention: A review in progress 1985-2005, and future research directions. Journal of Safety Research, 39, 259-267.
- CDC US Fall Prevention Programs for Seniors, <http://www.cdc.gov/ncipc/falls/fallprev.pdf>, accessed May 2008.
- CDC Injury Center, <http://www.cdc.gov/ncipc/factsheets/adultfalls.htm>, accessed April 2008.
- Rubenstein, LZ., Josephson, KR., Robbins, AS. Falls in the nursing home. Annals of Internal Medicine 1994;121:442-51.
- CDC Hip Fractures Among Older Adults, <http://www.cdc.gov/ncipc/factsheets/adulthipfx.htm>, accessed June 2008.
- Thomas, KE., Stevens, JA., Sarmiento, K., Wald, MM. (2008) Fall-related traumatic brain injury deaths and hospitalizations among older adults – United States, 2005. Journal of Safety Research, 39, 269-272.